

}vš šW > vv &}ÆU]Œ š}Œ }(À} Ç v Wµ o] ((]Œ•
o({Æ›]}%ZÇ•] •X}ŒP n ~îðì• îðirñòìò

& •š Œ D}vI Ç%}Æ ~u%}Æ• d •š]vP dZŒ}µPZ

Using gene editing proteins combined with nanopore sensing technology, new research shows that testing for pathogens could become faster and more accessible.

1
1

ROCKVILLE, MD Mpox, formerly known as monkeypox, is a rare viral disease that is spread 1 through physical contact between people. Currently, testing for mpox requires lab equipment and 1 can take a few hours to get test results. But new research suggests a way for faster testing that could 1 be done in any clinic soon. 1

1

Md. Ahasan Ahamed, a graduate student mentored by Weihua Guan at Pennsylvania State 1 University will present this research at the [68th Biophysical Society Annual Meeting](#), to be held 1 February 10-14, 2024 in Philadelphia, Pennsylvania. The work was also published in the February 1 issue of Biosensors and Bioelectronics. 1

1

Though mpox symptoms are generally mild with fever, rash, and swollen lymph nodes, severe cases 1 can occur and require medical attention. Because the disease is contagious, testing is important so 1 that people with the disease can isolate until symptoms resolve or get appropriate medical care. 1

1

To develop a faster test, the researchers used CRISPR, the Nobel prize winning gene editing 1 technology. Since 2017, scientists have expanded the application of CRISPR technology from gene 1 editing to molecular diagnostic techniques. 1

1


For this study, Ahamed created a genetic sequence combined with a reporter to specifically target 1 the mpox virus. Then a programmable CRISPR RNA binds to both the target and a protein called 1 Cas12a and together, the CRISPR/Cas12a leaves the reporter to create various sizes of fragments. 1 The researchers can then use nanopore sensing technology to analyze those reporters' fragments, 1 providing a rapid and accurate test that detects whether or not mpox is present in the sample. 1

1

The team confirmed that the test they created is specific to mpox—when they tested samples of 1 cowpox virus, a close relative of mpox, the test did not show a positive result. 1


1 1

The whole process is quick, "in total it takes 32 to 55 minutes to detect the target, depending on viral 1 load," Ahamed said, which is much faster than it currently takes to test load,"



The researchers plan to apply this nanopore technology to create tests for other pathogens, allowing one sample to be tested for multiple targets using portable device. And while the technology is not currently commercially available, Ahmed is hopeful that they will soon create a device that could make this kind of pathogen testing widely available.

1



1

1

CRISPR is combined with nanopore sensing technology to detect whether or not a pathogen is present in a sample. Image courtesy of Md. Ahasan Ahmed.

1

###

The Biophysical Society, founded in 1958, is a professional, scientific Society established to lead development and dissemination of knowledge in biophysics. The Society promotes growth in this expanding field through its annual meeting, publications, and committee and outreach activities. Its 7,500 members are located throughout the United States and the world, where they teach and conduct research in colleges, universities, laboratories, government agencies, and industry.